

## Drugs Used To Treat Infection: Antimicrobials



1



LESLIE MARTINEZ, MSNED, RN, MICN, PHN

LOS ANGELES COUNTY  
COLLEGE OF NURSING AND ALLIED HEALTH

LESLMARTINEZ@DHS.LACOUNTY.GOV

---

---

---

---

---

---

---

---



## What's the plan?

2

- **Antiinfectives**
  - Action/Indication
  - Administration
  - Safety: Labs
  - Side Effects
  - Adverse Effects
  - Patient Teaching
- **Drug Monitoring**
- **Food Considerations**
- **Nursing Process**
- **Drug Reactions**
- **Documentation**
- **Teaching**
- **Alternative Medicine**
- **Cultural Considerations**
- **NCLEX review game**

---

---

---

---

---

---

---

---

## Penicillins (PCN)

3



1. Pencillins
2. Broad-spectrum
3. Penicillinase-resistant
4. Extended-spectrum
5. Beta-lactamase Inhibitors
  1. Combination abx

📖 Table 29-3

---

---

---

---

---

---

---

---




---

---

---

---

---

---

---

---

## Basic Penicillins

5

- “Beta-lactam antibiotics”
- Interferes with & inhibits bacterial cell wall synthesis
- Narrow-spectrum
- Bacteriostatic & bacteriocidal
  - Dose dependent




---

---

---

---

---

---

---

---

## Basic Penicillins

6

- Indications of use
  - Staphylococcal infections
  - Severe wound & respiratory infections
- Examples
  - Penicillin G procaine
  - Bicillin




---

---

---

---

---

---

---

---

## Broad-spectrum PCNs

7

- “Aminopenicillins”
- Bactericidal
- Against gram + & gram –
- Indications of use: LRI, otitis media, sinusitis & UTI
- Examples: Amoxicillin & Ampicillin




---

---

---

---

---

---

---

## Broad-spectrum PCNs

8

- Amoxicillin
  - Most prescribed PCN derivative
  - Effect of Amoxicillin ↑ when taken with Aspirin & Probenecid
  - Effect of Amoxicillin ↓ when taken with Tetracycline & Erythromycin




---

---

---

---

---

---

---

## Amoxicillin

9

- Labs
  - Increased serum AST, ALT & BUN/Cr
- Food
  - Decreased effect with acidic fruits & juices
- Side effects
  - N/V/D, rash, edema
- Adverse reactions
  - Blood dyscrasias, hemolytic anemia, bone marrow depression, respiratory distress




---

---

---

---

---

---

---

## Penicillinase-resistant PCNs

10

- Effective against penicillinase-producing *Staphylococcus aureus*
- Against most gram +
  - *Less effective than PCN G*
- Examples
  - Dicloxacillin (PO) & Oxacillin (IM/IV)




---

---

---

---

---

---

---

---

## Extended Spectrum Penicillins

11

- Effective against *Pseudomonas aeruginosa*
- *Proteus* spp., *Serratia* spp., *Klebsiella pneumoniae*
- Broad Spectrum
- Indications of use
  - ✦ Treats bone, skin, respiratory tract infections & UTI
- Examples: Piperacillin & Ticarcillin




---

---

---

---

---

---

---

---

## Beta-lactamase Inhibitors

12

### Clavulanic acid, Sulbactam, & Tazobactam

- Indications for use
  - Penicillinase-producing *Staph. aureus*
- Combined with broad-spectrum abx
  - Extending the antimicrobial effect
- Adding Clavulanic Acid intensifies the effect of Amoxicillin (Augmentin)




---

---

---

---

---

---

---

---

## Beta-lactamase Inhibitors

13

- **PO: amoxicillin-clavulanic acid (Augmentin)**
  - Sinusitis, pneumonia & bronchitis
- **IM/IV: Piperacillin-tazobactam (Zosyn)**
  - UTI, bone & joint infections, stomach infections, skin infections & pneumonia
  - Reduce dose in renal insufficiency




---

---

---

---

---

---

---

## Drug Interactions

14



- Amoxicillin & ampicillin ↓ the effectiveness of oral contraceptives
- K<sup>+</sup> supplements ↑ serum K<sup>+</sup> levels when taken with Potassium PCN G or V
- PCN & Aminoglycosides mix in IVF → the actions of both are inactivated

---

---

---

---

---

---

---

## Safety: PCN

15

- **10% allergy rate: Monitor closely!**
- **Serum BUN/Cr, LFTs & Urine output**
  - Decrease dose with renal dysfunction
- ⚠ Side Effects
  - ⚠ Rash, itching, fever, chills, N/V/D
- ⚠ Adverse Reactions
  - ⚠ Hypersensitivity, superinfection, laryngeal edema, wheezing, hypotension




---

---

---

---

---

---

---

## Penicillins: Nursing Interventions

16

- C&S before med administration
- Monitor for bleeding
- Observe closely for allergic reaction
  - 1<sup>st</sup> and 2<sup>nd</sup> dose (epinephrine)
  - Medical Alert bracelet
- Increase fluid intake
- Assess for superinfection
  - Stomatitis & vaginitis




---

---

---

---

---

---

---

## Cephalosporins

17

- Four generations
- Beta-lactam structure
  - Semi-synthetic
- Inhibits bacterial cell-wall synthesis
  - Bactericidal
  - Cross-resistance with PCN
- Indications of use
  - Respiratory, urinary, skin, bone, joint & genital infections




---

---

---

---

---

---

---

## Cephalosporin Generations

18

### First-generation

- Gram + bacteria: *E. coli*, *Klebsiella*
- Cephalexin & cefazolin



### Second-generation

- Gram + & gram -: *Neisseria gonorrhoeae*, *Haemophilus influenzae*, *Neisseria meningitis*
- Cefaclor & cefoxitin

---

---

---

---

---

---

---

## Cephalosporins

19



- Third-generation
    - Gram + & gram -: *Pseudomonas aeruginosa*
    - Cefoperazone & ceftriaxone
  - Fourth-generation
    - Gram + & gram -: Streptococci, staphylococci
    - Cefepime
- 3 checks! Similar names.

---

---

---

---

---

---

---

---

## Cephalosporins

20



- Pharmacokinetics
  - Few PO; most IM, IV
- Side effects
  - N/V/D, GI distress & HA
- Adverse reactions
  - With high doses: Increased bleeding
  - Nephrotoxicity (in CRF/ARF)

---

---

---

---

---

---

---

---

## Cephalosporins

21

- Drug Interactions
  - **Alcohol:** may cause flushing, dizziness, HA, N/V & muscular cramps
  - **Oral Contraceptives:** decreases effect
  - **Uricosurics:** decrease Cephalosporin excretion (i.e. Probenecid)
    - ✦ Accumulation & toxicity




---

---

---

---

---

---

---

---

## Cephalosporins

22

- Nursing Interventions
  - Assess for allergic reaction
  - Assess renal & liver function
  - Monitor for superinfection
  - Monitor bleeding studies, VS & UO
- Safety
  - Keep out of reach of children




---

---

---

---

---

---

---

---

## Client Teaching

23

- Report s/s of superinfection
  - Instruct to ingest buttermilk or yogurt for prevention with long-term use
- Complete course of medication
- Childproof caps and out of reach of children
- Infuse all meds  $\geq 30$  min
- Report all side effects




---

---

---

---

---

---


---

---

## PCN Substitutes

24

- Macrolides, Lincosamides, Glycopeptides, & Ketolides
  - Broad-spectrum
  - Abx effectiveness similar to PCN
  - Used in PCN allergic patients




---

---

---

---

---

---

---

---



## Macrolides

25

- **Broad spectrum:** Some gram -/most gram +
  - Binds to ribosomal units & inhibits protein synthesis
  - Low to moderate doses: Bacteriostatic
  - High doses: Bacteriocidal
- **Indications of Use**
  - STIs, GI, respiratory, & skin infections
  - Commonly used in clients with PCN allergies




---

---

---

---

---

---

---

---

## Macrolides

26



- erythromycin (E-Mycin)
- **Extended Macrolides**
  - Longer half-life
  - azithromycin (Zithromax)
  - clarithromycin (Biaxin)

---

---

---

---

---

---

---

---

## Macrolides

27

- Increases effect of Digoxin, Tegretol, Theophylline, Cyclosporin & Warfarin
- **NOT** administered IM (OUCH!)
- **Azithromycin**
  - Decreases effect of PCNs & Clindamycin
  - Avoid antacids within 2 hours
- **Erythromycin**
  - Don't take with Clindamycin or Lincomycin




---

---

---

---

---

---

---

---

## Macrolides

28



- Safety: LFTs & UO

### Side Effects

- N/V/D & abdominal cramps

### Adverse Reactions

- Jaundice/icteric, anaphylaxis, superinfection, & hearing loss
- **Hepatotoxicity:** Erythromycin & Azithromycin + other hepatotoxics

---

---

---

---

---

---

---

## Nursing Interventions

29



- Obtain C&S
- Monitor VS & UO
- Monitor liver functioning:
  - Jaundice
  - LFT & bilirubin
- Instruct patient to report loose stools
  - Pseudomembranous colitis (C. dif)

---

---

---

---

---

---

---

## Lincosamides

30



- Inhibit bacterial protein synthesis
- Bacteriostatic & bacteriocidal
- Clindamycin (Cleocin)
  - Most gram +
- Side effects/Adverse reactions
  - N/V, stomatitis, colitis & anaphylactic shock

---

---

---

---

---

---

---




## Glycopeptides

31

- Glycopeptide bacteriocidal antibiotic

- Vancomycin (Vancocin)
- Telavancin (Vibativ) Once daily
  - ✦ Select gram + bacteria & skin infections
  - ✦ Effective against MRSA



- Indications of Use

- Cardiac surgical prophylaxis if PCN allergy

---

---

---

---

---

---

---

---

## Vancomycin

32

- Side Effects & Adverse Reactions

- Thrombophlebitis, N/V, dizziness, fever
- Ototoxicity & Nephrotoxicity

- Monitor

- Peak & trough levels
- Bun/Cr




---

---

---

---

---

---

---

---

## Redman Syndrome (Red Neck Syndrome)

33

- Red blotching of the face, neck & chest
- Toxic effect/Not an allergic reaction

- Treatment

- Mild to moderate:
  - ✦ Benadryl & Ranitidine, and ½ IV rate or 10mg/min
- Severe:
  - ✦ IVF PRN w/ hypotension
  - ✦ Administer Benadryl & Ranitidine
  - ✦ May restart once symptoms resolve (rate over 4 hours)

---

---

---

---

---

---

---

---

## Red Man Syndrome

34



---

---

---

---

---

---

---

## Stevens-Johnson Syndrome

35

- Epidermis separates from the dermis
- Hypersensitivity affecting skin & mucous membranes



---

---

---

---

---

---

---

## Vancomycin

36

- Risk of nephro & ototoxicity are potentiated with:
  - Furosemide, aminoglycosides, amphotericin B...
- Ototoxicity can be masked with Dramamine
- May inhibit methotrexate excretion = toxicity

---

---

---

---

---

---

---



## Ketolides

37

- Inhibits protein synthesis → bacterial cell death
- Telithromycin (Ketek)
  - >18 yo: Mild to mod. CA-pneumonia
- Side effects/Adverse reactions
  - HA, dizziness, N/V/D & **liver failure**

---

---

---

---

---



---

---

---

## Ketolide Drug Interactions

38

- Telithromycin levels  with:
  - Antilipidemics, itraconazole, ketoconazole & benzodiazepines
- Telithromycin levels  with:
  - Rifampin, phenytoin, carbamazepine & phenobarbital
- Increases multiple other drug levels

---

---

---

---

---

---


---

---

## Tetracyclines

39



- Broad-spectrum
  - Gram - & + & other organisms
  -  Bacterial resistance d/t overuse
  - Tx of H. Pylori → Peptic Ulcers w/ Metronidazole & bismuth
- Indications of use
  - Acne, rosacea, & **skin** infections
  - **Respiratory** infections & STIs

---

---

---

---

---

---

---

---

## Tetracyclines

(40)

- **Short-Acting**
  - tetracycline (Sumycin)
- **Intermediate-Acting**
  - demeclocycline (Declomycin)
- **Long-Acting**
  - doxycycline hyclate (Vibramycin)




---

---

---

---

---

---

---

---

## Tetracyclines

(41)

- **PO:** Rapid & complete absorption in newer preparations
- **IV:** Used to treat severe infections




---

---

---

---

---

---

---

---

## Tetracycline Administration

(42)

- Antacids, high calcium & iron drugs
  - Can prevent absorption of Tetracyclines
- Dairy products decrease Tetracycline effect
- **Doxycycline**
  - Decreases effect of oral contraceptives
  - Increases absorption of Digoxin → toxicity
  - Absorption enhanced with food




---

---

---

---

---

---

---

---

## Tetracyclines

45



### ⚠ Side Effects

- ⚠ N/V/D & photosensitivity
- ⚠ Teratogenic in 1<sup>st</sup> trimester (icon)

### ⚠ Adverse Reactions

- ⚠ Superinfections, hepatotoxicity, CNS toxicity...

⚠ Labs: BUN/Cr & LFTs

---

---

---

---

---

---

---

---

## Tetracycline Teeth

44




---

---

---

---

---

---

---

---


## Glycylcyclines

45

- Broad spectrum; gram + & -
- tigecycline (Tygacil)

### • Indications of Use

- Complicated skin infections, intrabdominal infections
- E. coli, S. aureus, Klebsiella pneumoniae




---

---

---

---

---


---

---

---

## Tigecycline

46

- Side effects & adverse reactions
  - N/V/D, **photosensitivity**, HTN/hypotension, anemia, leukocytosis & thrombocytopenia
  - Hyperglycemia, hypokalemia,  BUN & LFTs
- Concurrent therapy effects:
  - Oral contraceptives less effective
  - Warfarin levels may increase




---

---

---

---

---

---

---

---

## Aminoglycosides

47

- Inhibits bacterial protein synthesis
  - Gram -
  - Some gram + cocci are resistant → PCNs or Cephalosporins are used
- Indications of Use
  - PID, MRSA, & Pseudomonas
  - Serious infections




---

---

---

---

---

---

---

---

## Aminoglycosides

48

- amikacin (Amikin)
- gentamicin (Garamycin)
- neomycin (Mycifradin)
- tobramycin (Nebcin)
- No GI absorption: Primarily given IV & IM




---

---

---

---

---

---

---

---



## Aminoglycosides

49

### Side Effects

- N/V, rash & photosensitivity

### Adverse Reactions

- Ototoxicity, nephrotoxicity & liver damage**, thrombocytopenia & agranulocytosis

AMINOGLYCOSIDE TOXICITY



MAJOR TOXIC EFFECTS OF AMINOGLYCOSIDES ARE OTOTOXICITY &amp; NEPHROTOXICITY

## Aminoglycoside Safety

50

### Peak & Trough

- ↑ BUN, Serum AST, ALT, LDH, Cr & bilirubin
- ↓ Serum K<sup>+</sup> & Magnesium

#### 1. Nephrotoxicity

- Renal functioning, dose & age

#### 2. Ototoxicity risk ↑


- Concurrent ethacrynic acid treatment



## Nursing Interventions

51

- Check for hearing loss & balance
- Monitor UO
  - Report < 600 mL/24 hrs
- Peak and Trough
  - Gentamicin
- Monitor for superinfection
- Increase fluid intake



## Fluoroquinolones

52

- Broad Spectrum bacteriocidal
  - Interferes with enzyme DNA gyrase
  - Gram - & gram +
- Primary indications of use
  - UTI & Respiratory infections




---

---

---

---

---

---

---

---

## Fluoroquinolones

53

- ciprofloxacin HCl (Cipro)
- levofloxacin (Levaquin)
- ofloxacin (Floxin)
- moxifloxacin (Avelox)




---

---

---

---

---



---

---

---

## Fluoroquinolones Safety

54

- Food slows the absorption rate
-  Levofloxacin drug absorption with:
  - Antacids & Iron (within 4 hrs)
- Levofloxacin  effect of:
  - Oral hypoglycemics, Theophylline, & Caffeine
  - CNS reactions (tachycardia, anxiety...) can occur




---

---

---

---

---

---

---

---

## Fluoroquinolones

55

- Safety: BUN/Cr & UO
  - Elevated = kidney dysfunction
  - Fluid > 2,000 ml/day
- ☠ Side Effects
  - ☹ Dizziness, photosensitivity, N/V/D
- ☠ Adverse Reactions
  - ☹ Seizures, cardiac dysrhythmias & superinfection
  - ☹ Steven-Johnson syndrome




---

---

---

---

---

---

---

---

## Lipopeptides

56

- Bacteriocidal
- daptomycin (Cubicin)
- Indications
  - Complicated gram + skin infections, S. aureus septicemia and MRSA infective endocarditis




---

---

---

---

---

---


---

---

## Lipopeptides

57

- Side effects
  - HTN, hypotension & anemia
- Adverse reactions
  - Hyper/hypokalemia, hyper/hypoglycemia, bleeding & pleural effusion
- Toxicity with tobramycin
- Increased bleeding with Warfarin




---

---

---

---

---

---

---

---



## Sulfonamides

58



- Bacteriostatic
  - Inhibit bacterial synthesis of folic acid
  - Gram –
  - Prevents bacterial growth in the kidneys & bladder
- Indications of Use
  - UTI, ear infections, meningococcal meningitis, newborn eye prophylaxis & STIs

---

---

---

---

---

---

---

---

## Sulfonamides

59

- Short-Acting
  - sulfadiazine (Microsulfon)
- Intermediate-Acting
  - trimethoprim-sulfamethoxazole (Bactrim, Septra)
- Topical
  - silver sulfadiazine (Silvadene)
- Can be used in clients with PCN allergy

---

---

---

---

---

---

---

---

## Sulfonamides

60



- ⚠ Side Effects/Adverse Reactions
  - ⚠ Skin rash/itching, photosensitivity, cross-sensitivity, & N/V/D
- ⚠ Prolonged use
  - ⚠ Hemolytic or aplastic anemia, thrombocytopenia, neutropenia, & agranulocytosis

---

---

---

---

---

---

---

---

## Nursing Interventions

61

### Teaching

- ✍ Crystalluria → Increase fluid intake
- ✍ Sunglasses, report bruising or bleeding
- Assess renal functioning
  - BUN/Cr and UO
  - Bactrim and Septra: contraindicated in renal clients
- **Increases hypoglycemic effect with hypoglycemics**
- **Increases anticoagulant effect with Warfarin**




---

---

---

---

---

---

---

## Urinary Tract Disorder Drugs

62

- Upper UTI
  - Pyelonephritis
- Lower UTI
  - Cystitis, urethritis or prostatitis
- UA and C&S prior to tx
- Acute Cystitis
  - Demographics
  - Signs & Symptoms




---

---

---

---

---

---

---

## Definitions

63

- Urinary Antiseptics/antiinfectives:
  - Prevents bacterial growth in the kidneys & bladder
    - ✦ Sulfonamides/Fluoroquinolones
- Urinary Analgesics:
  - Relieve urinary tract pain & burning
- Urinary Stimulants:
  - Increase urinary muscle tone




---

---

---

---

---

---

---

### Urinary Antiseptics/Anti-infectives

64

- nitrofurantoin (Macrochantin)
  - Bacteriostatic & Bacteriocidal (dose dep.)
  - Gram - & +
- Indications of Use
  - Acute & Chronic UTI




---

---

---

---

---

---

---

---

### Urinary Antiseptics/Anti-infectives & Antibiotics

65

- nitrofurantoin (Macrochantin)
- Adverse Reactions
  - Dyspnea, chest pain, fever/chills
- Nursing Considerations
  - ↓ absorption with antacids
  - Accumulates in serum with urinary dysfunction

---

---

---

---

---

---

---

---

### Urinary Antiseptics/Anti-infectives & Antibiotics

66

- methenamine (Hiprex)
  - Bacteriocidal when pH < 5.5
- Indications of Use
  - Chronic UTI
- Nursing Considerations
  - Cystalluria occurs with sulfonamides
  - Encourage acidic juices & ascorbic acid (vit c)
  - Decrease alkaline food intake




---

---

---

---

---

---


---

---

## Urinary Analgesics

67

- pyridium (Phenazopyridine)
- Urinary Analgesic: Azo dye
  - Relieves pain, burning, frequency & urgency
- Indications of Use
  - Lower UTI
  - Concurrent antibiotic tx




---

---

---

---

---

---


---

---

## Pyridium (Phenazopyridine)

68

- Clinitest: Alters the result
- ⚠ Side effects/ Adverse reactions
  - ⚠ GI disturbances, hemolytic anemia, blood dyscrasias, nephrotoxicity & hepatotoxicity
- Teaching
  - Reddish-orange urine from dye
  - May stain contact lenses
  - Report N/V/D




---

---

---

---

---

---


---

---

## Urinary Stimulants

69

- Parasympathomimetics
- Urecholine
  - ↑ bladder tone
  - Produces contraction that stimulates micturition
- Which type of client would benefit from this drug class?




---

---

---

---

---

---

---

---

## Urinary Antispasmodics/Antimuscarinics

70

- Relieve urinary tract spasms from infection
- dimethyl sulfoxide, oxybutynin & flavoxate
- Contraindications
  - GI obstruction
  - Glaucoma




---

---

---

---

---

---

---

---

## Urinary Antispasmodics/Antimuscarinics

71

- Side Effects:
  - Dry mouth, ↑ HR, dizziness, GI distention & constipation
- Teaching
  - Report retention, severe dizziness, blurred vision, palpitations & confusion
  - Avoid prolonged heat exposure

---

---

---

---

---

---

---

---

## Adverse Effects of Antibacterials

72

- Allergy or hypersensitivity reaction
  - Mild: Rash, pruritis & hives → antihistamine
  - Severe: Laryngeal edema, bronchospasms, cardiac arrest → epi, bronchodilator & antihistamine
  - Usually occurs in the first 20 minutes
- Superinfection
- Organ toxicity
  - Liver & Kidneys




---

---

---

---

---

---

---

---



## Superinfection

75

- 📖 Secondary infection: when normal flora are killed
- 📖 Sites: Mouth, skin, resp. tract, vagina, intestines
- 📖 Fungal or Bacterial
- 📖 Broad-spectrum > 1 wk

### 📖 Adverse reaction

- 📖 Notify the MD stat!



©ADAM

---

---

---


---

---

---

---

---




---

---

---

---

---

---


---

---

## Resistance

75

- ☑️ Lessened antibiotic effect d/t misuse
  - ☑️ Used on viruses or wrong bacteria/fungus
  - ☑️ Mutant bacteria survive antibiotic use
  - ☑️ Taken incorrectly
- ☑️ Combating resistance
  - ☑️ New classes of drugs
    - ☑️ Abx resistant disabler
  - ☑️ Bacterial vaccines (pneumococcal)
  - ☑️ Patient teaching




---

---

---

---

---


---

---

---

## Peak & Trough Review

76



- Peak
  - Drawn @ drug's peak of action
  - Indicates rate of drug absorption
- Trough
  - Drawn minutes before drug administration
  - Indicates rate of drug elimination
- Gentamicin Peak 5-10      Toxic Peak > 12  
                                  Trough < 2      Toxic Trough > 2  
     ~Peak is 30 minutes after IV administration

---

---

---

---

---

---

---

---

## Culture & Sensitivity (C&S)

77



- 📖 Detects the infective microorganism in the blood & what drug can kill it
- 📖 **Culture:** Organism causing the infection
- 📖 **Sensitivity:** Antimicrobial the organism is sensitive to
- 📖 Draw **before** antimicrobial administration

---

---

---

---

---

---

---

---



## Liver & Kidney Tests

78



- CLcr
  - **Most accurate** lab test in determining **renal function**
  - Determines dose adjusting
- BUN
  - Determines renal functioning or dehydration (high levels)
- Creatinine
  - Specific indicator of renal functioning
- LFTs
  - ALT
  - AST
  - Alk Phos




---

---

---

---

---

---

---

---

## Dose Adjusting


79

### • Organ Toxicity

- Liver, Kidney & Ears
- CLcr, BUN/Cr & LFT's
- Jaundice, UO, auditory checks

### • Critically-ill patients

- Immunocompromised




---

---

---

---

---

---

---

---

## Nursing Process: Assessment

80

### • Allergies

### • Lab Results

- LFTs, Peak & Trough, C&S

### • Record/Report UO, VS

### • Diet

### • History

### • Physical Assessment




---

---

---

---

---

---

---

---

## Physical Assessment Findings

81



- Describe wound, surgical site...

- Monitor for allergic reaction, notify MD & document findings **promptly**

- Rash, wheezing, fever...

- Kidney/Liver damage

- Jaundice/Icteric

- Urine output decreased

- Superinfection



Hives

#ADAM

81

---

---

---

---

---

---

---

---

## Nursing Diagnosis

82

1. Risk for impaired tissue integrity r/t rash.
2. Risk for infection r/t invasion of bacteria through surgical incision.
3. Noncompliance with drug regimen r/t lack of knowledge relevant to regimen behavior AEB taking two days of 11 day therapy.

---

---

---

---

---

---

---

## Planning/Goals

83

### • Short Term

- Client will demonstrate meticulous hand washing technique by 1300.
- Client will demonstrate knowledge of risk factors associated with infection by 1100.




---

---

---

---

---

---

---

## Interventions

84

- Send labs, swabs...
  - **Before** antibiotic administration
- Assess for Allergic Reaction
  - Medical Alert bracelet
- Monitor temperature
  - 1 hr after antipyretic
  - Administer 2<sup>nd</sup> antipyretic prn




---

---

---

---

---

---

---

## Patient Teaching

85



- Teaching Plan
  - Indication of use
  - Schedule/duration
  - Food considerations
  - Interaction with herbs & other medications
- Validation of Understanding
- STI Prevention
- Hand Washing
- Take full course of antibiotic as directed
  - Finish full Rx
  - Prevent resistance
- Report side effects/adverse reactions
  - Superinfection

---

---

---

---

---

---

---

## Evaluation

86

- Infection ceased
- Adverse reactions
  - Superinfection
- WBC count WNL
- Afebrile
- VSS: i.e. HR, RR




---

---

---

---

---

---

---

## Cultural Considerations

87

- Alternative Practices
  - Curandero/healer
- Language Barrier
  - Translator
- Drug Schedule
  - Alternative methodology




---

---

---

---

---

---

---

## Complimentary, Alternative & Traditional Medications

88

### • Client Preference

- Homeopathic vs. Rx
  - Interactions with Rx
- Previous experience



### • Teaching

- Stress importance of medical regime

### • Family Involvement

- Teach client with family at bedside

---

---

---

---

---

---

---

---

## Questions?

© Original Artist  
Reproduction rights obtainable from  
www.CartoonStock.com



"It's a prescription for one of those new super-antibiotics. You won't just get better, you'll get even."

---

---

---

---

---

---

---

---